

# Factors influencing post-exercise plasma protein carbonyl concentration

Aldred, Sarah; Turner, James; Wadley, Alex

DOI:

[10.3109/10715762.2015.1131824](https://doi.org/10.3109/10715762.2015.1131824)

License:

Other (please specify with Rights Statement)

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Aldred, S, Turner, J & Wadley, A 2016, 'Factors influencing post-exercise plasma protein carbonyl concentration', *Free Radical Research*, vol. 50, no. 4, pp. 375-384.  
<https://doi.org/10.3109/10715762.2015.1131824>

[Link to publication on Research at Birmingham portal](#)

## **Publisher Rights Statement:**

This is an Accepted Manuscript of an article published by Taylor & Francis in *Free Radical Research* on 12/02/2016 available online:  
<http://www.tandfonline.com/10.3109/10715762.2015.1131824>

## **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## **Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

# **Factors influencing post-exercise plasma protein carbonyl concentration**

*Alex, J. Wadley<sup>1</sup> James, E. Turner<sup>2</sup> & Sarah Aldred<sup>3</sup>*

<sup>1</sup> Institute of Science and the Environment, University of Worcester, Worcester, WR2 AJ

<sup>2</sup> School of Sport, Exercise & Rehabilitation Sciences, University of Birmingham, Edgbaston, Birmingham, B15 2TT

<sup>2</sup> Department for Health, University of Bath, Bath, BA2 7AY, UK.

**Running Head:** Protein carbonyl groups and exercise

**Word Count:** 2967

## **Address for correspondence:**

Dr Sarah Aldred

School of Sport, Exercise and Rehabilitation Sciences

College of Life & Environmental Sciences

University of Birmingham

B15 2TT, United Kingdom

Email: s.alred.1@bham.ac.uk

Phone: 0121 414 7284

Fax: 0121 414 4121

## **Abstract**

Exercise of sufficient intensity and duration can cause acute oxidative stress. Plasma protein carbonyl (PC) moieties are abundant, chemically stable and easily detectable markers of oxidative stress that are widely used for the interpretation of exercise-induced changes in redox balance. Despite many studies reporting acute increases in plasma PC concentration in response to exercise, some studies, including those from our own laboratory have shown decreases. This review will discuss the differences between studies reporting increases, decreases and no change in plasma PC concentration following exercise in humans; highlighting participant physiology (i.e. training status) and study design (i.e. intensity, duration and novelty of the exercise bout) as the main factors driving the direction of the PC response to exercise. The role of the 20S proteasome system is proposed as a possible mechanism mediating the clearance of plasma PC following exercise. Resting and exercise-induced differences in plasma protein composition and balance between tissues are also discussed. We suggest that exercise may stimulate the clearance of plasma PC present at baseline, while simultaneously increasing reactive oxygen species production that facilitates the formation of new PC groups. The balance between these two processes likely explains why some studies have reported no change or even decreases in plasma PC level post-exercise when other biomarkers of oxidative stress (e.g., markers of lipid peroxidation) were elevated. Future studies should determine factors that influence the balance between PC clearance and formation following acute exercise.

**Keywords:** Protein oxidation, exercise, proteasome, protein degradation, reactive oxygen species

## **Introduction**

Exercise can induce a wide range of whole body physiological adaptations that improve metabolic health and lower oxidative stress [1–3]. Oxidative stress is a biological state whereby reactive oxygen species (ROS) overwhelm antioxidant defences, increasing the oxidation of proteins, lipids and DNA. It is widely accepted that transient increases in exercise-induced ROS can initiate a diverse range of signalling pathways that lead to adaptation [4–6]. Indirect biomarkers of exercise-induced oxidative stress, such as protein oxidation [7–9], lipid peroxidation [8–10] and antioxidant capacity [8,11,12] are routinely measured to give an indication of altered redox balance. One of the most frequently examined biomarkers of protein oxidation is plasma protein carbonyl (PC) concentration. Carbonylation is a stable and quantifiable post-translational protein modification which is ten times more abundant than other protein adducts, such as 4-Hydroxynonenal and glycooxidation end-products [13,14]. Most biomarkers of oxidative stress increase in plasma in response to exercise [11,12,15], as would be expected following an acute bout of increased metabolic activity. However some studies have reported decreases in plasma PC concentration post-exercise alongside increases in other biomarkers of oxidative stress [8,16,17]. The focus of this review is to explore key physiological factors that might explain these different responses, with a primary emphasis on aerobic, steady state exercise bouts where PC groups have been measured in blood plasma or serum of human participants exercising under fasted conditions. It is beyond the scope of this review to include the results of studies that have examined the effect of habitual diet or dietary supplementation (e.g., high dose antioxidants) on resting or exercise-induced changes in PC levels.

## **Protein carbonyl formation**

PC groups are present in all proteins (carboxylic acid (-COOH) groups), form the basis of their structural integrity, and influence their capacity to function and interact with other molecules. Formation of additional, non-native PC groups can be a result of a variety of irreversible, non-enzymatic oxidative pathways (*Figure 1*) that are a normal part of metabolic processes [18]. These include direct oxidation of amino acids (in conjunction with oxidising agents such as transition metal clusters in protein structures) and secondary oxidation reactions with lipid peroxidation or glucose-protein oxidation products [19]. PC groups are highly polar, thus increasing the proteins susceptibility to further oxidation and/or formation of cross-linkages and protein aggregates (via ketone or aldehyde links) [20]. Excessive introduction of new PC groups can result in altered or disrupted protein function and exposure of previously embedded hydrophobic groups in the protein core, resulting in targeted proteolytic degradation by 20S and 11S proteasome systems [20,21]. The degree of PC formation is dependent on the presence of conjugated metals (e.g., iron [19]), the orientation of specific amino acids that are more susceptible to PC formation (i.e., Proline, Arginine, Lysine, and Threonine) and importantly, the magnitude and origin of ROS production in relation to the protein [22].

*[Insert Figure 1 here]*

### **Exercise-induced changes in protein carbonylation**

A variety of cells can produce ROS in response to exercise, most notably myocytes [23], leukocytes [24] and endothelial cells [25]. Superoxide ( $O_2^{\cdot-}$ ) is produced from a range of enzymatic sources within these cell types during exercise, such as nicotinamide adenine dinucleotide phosphate (NADPH) oxidase, xanthine oxidase and nitric oxide synthases [26]. These cells can also release  $O_2^{\cdot-}$  into the extracellular space via enzymes expressed on the

plasma membrane such as NADPH oxidase [26–28] or via passive diffusion of uncharged ROS, such as hydrogen peroxide [29]. Consequently, plasma proteins are susceptible to the formation of PC groups during exercise [7,9,30,31], with evidence that new PC moieties are stable in plasma for up to 4 hours, before selective degradation [32,33]. Increased formation of plasma PC groups following exercise is considered to be a non-specific reflection of increased systemic oxidative stress (i.e. the origin of ROS and biological impact is unknown). As a result, many studies over the last 15 years have investigated associations between changes in plasma PC concentration and aspects of the acute physiological stress caused by exercise, in a variety of populations [7,9].

## **Factors influencing post-exercise changes in protein carbonylation**

### **Exercise intensity**

Aside from the potential for greater production of ROS from the cellular sources discussed above, specific hypoxic mechanisms can also contribute during high intensity exercise. Repetitive cycles of temporary occlusion (hypoxia) and re-oxygenation of the blood vessels surrounding actively contracting muscle can produce large quantities of ROS via the enzyme xanthine oxidase [34], that may increase PC formation. Previous studies have reported increases in plasma PC concentration following high-intensity exercise to exhaustion relative to baseline values [12,31,35–37]. Lamprecht et al [30] assessed the impact of exercise intensity on plasma PC formation by examining three 40-minute cycling bouts of different intensities (70, 75 and 80%  $\dot{V}O_{2\max}$ ) in three independent groups of moderately active participants (*Table 1*). Plasma PC concentration increased in response to cycling at 80%  $\dot{V}O_{2\max}$  only, suggesting that exercise intensity is a determinant of the formation of new non-native carbonyl moieties in the bloodstream. However it is likely that exercise duration and other physiological factors also have an impact upon these processes.

75

76 Exercise duration

77       There is some evidence to suggest that exercise duration is also a key factor in post-  
78 exercise plasma PC formation. Bloomer et al [7] reported that 120 minutes of cycling at 70%  
79  $\dot{V}O_{2\max}$  caused a greater increase in post-exercise plasma PC concentrations than 30 and 60-  
80 minute bouts of the same exercise intensity in male and female participants. Moreover, an  
81 exercise bout of moderate intensity, but long duration (ultra-endurance running: 174 km, 30-  
82 44 hours) has been reported to elicit immediate and prolonged (7 days) post-exercise  
83 increases in plasma PC concentration [38]. This increased protein oxidation occurred  
84 simultaneously with a decline of exogenous and/or endogenous antioxidants, which may have  
85 reduced the capacity to clear PC groups within this seven day period. The roles of exercise  
86 intensity and duration on post-exercise PC concentrations are inevitably linked; however  
87 there is evidence that other physiological factors (e.g., training status) contribute [7,15].

88

89 Training status and habituation to exercise

90       Exercise-induced ROS production can initiate a cascade of cellular signals which  
91 result in various post-translational modifications (i.e. phosphorylation, acetylation and thiol  
92 modifications), and up-regulate the expression of antioxidant and stress proteins following  
93 exercise [4,39,40]. Differences in the resting expression of antioxidant proteins between  
94 participants (i.e. due to training status and/or habituation to exercise) will no doubt have  
95 consequences for changes in exercise-induced oxidative stress. Indeed, it has been  
96 demonstrated that exercise training can stimulate an increased expression of endogenous  
97 antioxidant proteins [5,41], with some direct evidence of this in humans [42]. However, it is  
98 largely unclear how much resting antioxidant capacity impacts upon the acute oxidative  
99 stress response to exercise. Elevated endogenous antioxidant capacity may enable a buffering

of ROS production subsequently reducing the magnitude of oxidative stress biomarker formation during exercise in trained individuals [43]. Bloomer et al [7] reported increases in plasma PC concentration immediately following cycling exercise (30 minutes at 70%  $\dot{V}O_{2max}$ ) (Table 1), but in a separate study, with different research participants, reported no change in plasma PC level following an identical cycling protocol [15]. Differences in the physiology and exercise training habits of the participants featured in these studies, rather than the intensity and duration of the exercise bout are likely to have influenced the differential net changes in PC post-exercise. For example, there were clear differences in both the aerobic fitness ( $\dot{V}O_{2max}$ :  $57 \pm 5$  ml/kg/min [7] vs.  $45 \pm 8$  ml/kg/min [15]) and time engaged in aerobic exercise prior to the study ( $10.4 \pm 1.6$  hours/week [7] vs.  $2.8 \pm 2.2$  hours/week [15]). In addition, the participants in the study published by Bloomer et al, 2005 [15] were more resistance ( $3.8 \pm 1.8$  hours/week), than aerobically trained. This suggests that the exercise stimulus implemented in both studies was more novel for the participants in the 2005 study [15]. This supports some previous work reporting a greater magnitude of oxidative stress biomarker formation following unaccustomed exercise [44,45]. Thus, a combination of factors such as the novelty of exercise, and aspects of study design (i.e., exercise intensity and duration) are likely to interact to govern the magnitude of increase in plasma PC formation following exercise. However, it is perhaps more challenging to explain why *decreases* in plasma PC level have been observed.

#### **Studies reporting *decreases* in plasma PC concentration after exercise**

The available evidence suggests exercise intensity [30] and duration [7] influence the magnitude of ROS production and the associated increase in biomarkers of oxidative stress. It is therefore not surprising that many studies have reported no change in plasma PC concentration following sub-maximal exercise [15,46–50]. However, counter-intuitively,



many studies report a *decrease* in plasma PC concentration following both sub-maximal and maximal exercise [8,16,17,51] (*Table 1*). Importantly, these changes have sometimes been reported alongside increases in other biomarkers of oxidative stress [8,17]. This is unexpected and implies that exercise stimulates clearance processes alongside exercise-induced ROS production.

We have shown that steady state submaximal cycling (60% and 80%  $\dot{V}O_{2max}$  for 27 [moderate intensity] and 20 minutes [high intensity] respectively) undertaken by untrained males ( $\dot{V}O_{2max}$ ;  $42.7 \pm 5.0$  ml/kg/min) elicits a 9% (moderate intensity,  $p < .0001$ ) and 4% (high intensity,  $p < .0001$ ) mean decrease in plasma PC concentration (see figure 2: A and B) [8]. Importantly, these changes occurred in parallel with increases in plasma lipid hydroperoxides (LOOH), total antioxidant capacity (TAC) and cellular markers of oxidative stress [52]. Interestingly, the participants in this study engaged in less than 3 hours of generic aerobic exercise per week, indicating that these bouts were relatively unaccustomed. These findings are not limited to just moderately trained individuals. We have also shown a 7% decrease ( $p = .016$ ) in plasma PC concentration immediately after a bout of cycling exercise (70%  $\dot{V}O_{2max}$ , 75 min) in highly trained cyclists ( $\dot{V}O_{2max}$ ;  $63.7 \pm 5.3$  ml/kg/min) (Wadley et al, 2015; In Preparation; see figure 2D). Finally, our findings are not limited to sub-maximal exercise: we have also shown a 10% mean decrease ( $p = .002$ ) in plasma PC concentration immediately after a graded exercise test to volitional exhaustion (i.e., 100%  $\dot{V}O_{2max}$ , approximately 15-minutes) in very active young men ( $\dot{V}O_{2max}$ ;  $61.9 \pm 4.7$  ml/kg/min; see Figure 2C) (Turner *et al*; unpublished data) and a 13% mean decrease ( $p < .0001$ ) following a bout of low volume high intensity interval exercise ( $10 \times 1$  minute stages at 90%  $\dot{V}O_{2max}$ , with  $9 \times 1$  minute rest intervals at 40%  $\dot{V}O_{2max}$ ) in moderately active participants ( $\dot{V}O_{2max}$ ;  $42.7 \pm 5.0$  ml/kg/min) [8].

There are a number of other studies that report (but do not always comment upon) decreases in plasma PC levels following exercise (Table 1). For example, Chevion et al, [16] reported decreases in PC concentration following a study involving two walks (50km and 80km) of unreported intensity. The magnitude of this decrease in PC concentration was greater following the first walk compared to the second, highlighting again that the novelty of the exercise stimulus may be a key factor modulating post-exercise PC concentration. Furthermore, reductions in plasma PC level have been reported immediately [17,51] and up to four hours [17] following submaximal cycling exercise in moderately trained participants.

*[Insert Figure 2 here]*

## **Possible mechanisms mediating protein carbonyl clearance in response to exercise**

### **The 20S proteasome system**

The proteasome system is an organised assembly of proteins present in all cell types that functions to degrade irreversibly modified proteins, such as those containing carbonyl groups. The ubiquitin-independent 20S proteasome is the primary system in place to degrade oxidatively damaged proteins [21,32,53], with recent evidence also suggesting the 11S proteasome facilitates this process under conditions of heightened oxidative stress [20]. Increased exposure of carbonyl-mediated hydrophobic groups within the protein core can increase targeted degradation by these proteasome systems within cells. The 20S proteasome is also excreted into extracellular fluids such as plasma [54–56] and is known to be enzymatically functional [57], suggesting that it could cleave PC groups in plasma directly. Studies in humans have found that exercise can acutely increase ubiquitin-dependent proteasome gene expression in response to resistance [58–60] and ultra-endurance exercise [61]. However, the impact of acute shorter-duration aerobic-type exercise on the ubiquitin-

independent 20S and 11S proteasome subunits (either intra- or extra-cellular; expression or activity) and the relationship with plasma PC has not been investigated in humans. Data from rats has indicated that chymotrypsin-like activity of the ubiquitin-independent 20S proteasome subunit increases in brain tissue following 8 weeks of exercise overload (1 hour swimming/day, 5 days a week for 6 weeks) [62].

Exercise bouts below a certain intensity and/or duration, in combination with the other physiological parameters discussed above may activate intracellular and extracellular proteasome pathways to clear modified proteins and thus lower the concentration of PC groups in plasma (*Figure 3*). The net increase in PC level observed in response to high intensity or prolonged duration exercise may result from the formation of new PC groups outnumbering proteolytic clearance of those present at baseline. Furthermore, inferences from cell culture data [53] led Radak et al [63] to propose that proteasome activity may follow a hormetic-type response with increasing exercise-induced oxidative stress. Proteasome activity may be reduced at higher exercise intensities or after prolonged duration exercise, due to ROS-induced inactivation of the functional 20S or 11S proteasome. Future work is needed to explore this mechanism.

#### Resting or exercise-induced differences in total plasma protein composition and balance between tissues

Studies examining plasma PC concentration typically express results relative to the total protein concentration of plasma (i.e., nM of carbonyls per mg of protein) [64]. Measuring protein carbonylation in this way does not account for possible exercise-induced shifts in protein composition. This is relevant because certain plasma proteins, such as serotransferrin and fibrinogen (approximately 2-4% of total plasma proteins) are more susceptible to exercise-induced oxidation than other plasma proteins [18]. Therefore,

individual differences in baseline plasma protein composition may influence post-exercise changes in PC. Furthermore, exercise could induce shifts in the proportion of certain proteins in plasma, which could result in exaggerated or suppressed protein carbonylation.

Although not as susceptible to oxidation as fibrinogen, albumin, the most abundant plasma protein, has also been shown to exhibit a high concentration of PC groups [65]. The extent to which albumin becomes oxidised in response to exercise is dependent on intensity, and albumin carbonylation is marked at intensities of 80%  $\dot{V}O_{2\max}$  or more [30]. Inter-individual variation in exercise-induced proteinuria or albuminuria has been documented [66,67], and might affect the amount and type of plasma proteins that can be oxidised during exercise. Furthermore, it has been shown that during and following both aerobic and resistance exercise, there is an increase in the uptake and turnover of proteins such as albumin and fibrinogen by muscle [68,69]. Thus, differences in PC excretion and balance between tissues might explain to some degree, alterations in the composition of plasma proteins, which may increase or decrease plasma PC level.

#### **Experimental approaches for monitoring changes in plasma protein carbonyl level**

The evidence presented in this review highlights the uncertainty with regards to the mechanistic underpinnings of post-exercise decreases in plasma protein carbonylation. Despite the factors discussed, experimental design and the analytical techniques used to quantify protein carbonyl concentration in the cited studies warrants discussion.

The majority of studies included in *Table one* controlled for factors such as age, training status, dietary habits (i.e. fasted exercise trials), and the time of day that exercise was undertaken. All of these variables have been shown to alter protein balance through changes in protein uptake to skeletal muscle during exercise [68,69], highlighting the rigorous experimental approach taken in these studies. However, an important consideration is the

224 timing of blood sampling. Many of the studies presented assessed protein carbonyl  
225 concentration immediately following cessation of the exercise bout. It is conceivable that  
226 changes within the minutes and hours after exercise might give more insight into the  
227 ‘conundrum’ of post-exercise changes in protein carbonyl level. Indeed, independent studies  
228 have shown increases [70] and decreases [17] in protein carbonyl concentration up to 4 hours  
229 post-exercise. In this regard, it is clear that extensive timecourse analysis of protein carbonyl  
230 concentration is needed to validate these findings and importantly, to elucidate the  
231 mechanisms influencing decreases following exercise.

232         It is important to note that a variety of analytical techniques are used to quantify  
233 protein carbonylation in the literature. These have primarily included spectrophotometric and  
234 ELISA-based methods, which have recently come under scrutiny with regards to their  
235 sensitivity [71] and reproducibility [72]. However, while these studies provide an example of  
236 an important limitation of quantifying protein carbonyl concentration, they do not explain  
237 why protein carbonyl concentration might decrease following exercise. Furthermore, with  
238 independent laboratories now reporting decreases in protein carbonyl level following exercise  
239 [8,16,17,51], this reduces the chances of experimental variability causing this effect.

240  
241 *[Insert Figure 3 here]*  
242

## 243 **Conclusions**

244         Plasma protein carbonyl concentration is a marker of oxidative stress routinely used  
245 to assess exercise-induced redox regulation. The evidence presented in this review suggests  
246 that certain exercise conditions can result in a net *decrease* in plasma PC concentration  
247 following exercise, which occurs in parallel with increases in other biomarkers of oxidative  
248 stress. Exercise intensity ( $>70\% \dot{V}O_{2\max}$ ) and prolonged duration ( $>60$  minutes) appear to be

the main contributing factors in the observed post-exercise increases in PC concentration. The factors influencing *decreases* in protein carbonyl level are more difficult to interpret, but likely involve the clearance of oxidised proteins from plasma, potentially by plasma proteasomes, excretion, or uptake into active tissues. Studies wishing to assess markers of oxidative stress in response to exercise should assess PC concentration together with other biomarkers over an extended time course (immediately after and up to 4 hours post-exercise) for a true representative assessment [31]).

## Acknowledgments

All authors contributed to the following aspects of this study: conception and outline of the brief review, collection and analysis of the unpublished data presented, and drafting the article and revising it critically for important intellectual content.

## Declaration of Interest

None of the authors declare a conflict of interest and have no financial interest in the study.

## References

- [1] Fatouros IG, Jamurtas AZ, Villiotou V, Pouliopoulou S, Fotinakis P, Taxildaris K, Deliconstantinos G. Oxidative Stress Responses in Older Men during Endurance Training and Detraining. *Med Sci Sport Exerc* 2004;2065–2072.
- [2] Donato AJ, Uberoi A, Bailey DM, Wray DW, Richardson RS. Exercise-induced brachial artery vasodilation: effects of antioxidants and exercise training in elderly men. *Am J Physiol Hear Circ Physiol* 2010;298:H671–8.
- [3] Radak Z, Kaneko T, Tahara S, Nakamoto H, Ohno H, Sasvari M, Nyakas C, Goto S. The effect of exercise training on oxidative damage of lipids, proteins, and DNA in rat skeletal muscle: evidence for beneficial outcomes. *Free Radic Biol Med* 1999;27:69–74.
- [4] Gomez-Cabrera MC, Domenech E, Romagnoli M, Arduini A, Borrás C, Pallardo F V., Sastre J, Viña J. Oral administration of vitamin C decreases muscle mitochondrial biogenesis and hampers training-induced adaptations in endurance performance. *Am J Clin Nutr* 2008;87:142–9.

- 280 [5] Gomez-Cabrera MC, Domenech E, Vina J. Moderate exercise is an antioxidant:  
281 upregulation of antioxidant genes by training. *Free Radic Biol Med* 2008;44:126–131.
- 282 [6] Ristow M, Birringer M, Kiehntopf M, Zarse K, Oberbach A, Klo N, Stumvoll M,  
283 Kahn CR, Blu M. Antioxidants prevent health-promoting effects. *PNAS*  
284 2009;106:8865–8670.
- 285 [7] Bloomer RJ, Davis PG, Consitt LA, Wideman L. Plasma protein carbonyl response to  
286 increasing exercise duration in aerobically trained men and women. *Int J Sports Med*  
287 2007;28:21–25.
- 288 [8] Wadley AJ, Chen YW, Lip GY., Fisher JP, Aldred S. Low volume-high intensity  
289 interval exercise elicits anti-oxidant and anti-inflammatory effects in humans. *J Sports*  
290 *Sci* 2015;27:1–9.
- 291 [9] Wadley AJ, Veldhuijzen van Zanten JJCS, Stavropoulos-Kalinoglou A, Metsios GS,  
292 Smith JP, Kitas GD, Aldred S. Three months of moderate-intensity exercise reduced  
293 plasma 3-nitrotyrosine in rheumatoid arthritis patients. *Eur J Appl Physiol*  
294 2014;114:1483–92.
- 295 [10] Wadley AJ, Veldhuijzen van Zanten JJCS, Paine NJ, Drayson MT, Aldred S.  
296 Underlying inflammation has no impact on the oxidative stress response to acute  
297 mental stress. *Brain Behav Immun* 2014;40:182–90.
- 298 [11] Berzosa C, Cebrian I, Fuentes-Broto L, Gomez-Trullen E, Piedrafita E, Martinez-  
299 Ballarin E, Lopez-Pingarron L, Reiter RJ, Garcia JJ. Acute exercise increases plasma  
300 total antioxidant status and antioxidant enzyme activities in untrained men. *J Biomed*  
301 *Biotechnol* 2011;1–7.
- 302 [12] Wadley AJ, Killer SC, Svendsen IS, Gleeson M. The impact of intensified training  
303 with a high or moderate carbohydrate feeding strategy on resting and exercise-induced  
304 oxidative stress. *Eur J Appl Physiol* 2015;
- 305 [13] Jung T, Höhn A, Catalgol B, Grune T. Age-related differences in oxidative protein-  
306 damage in young and senescent fibroblasts. *Arch Biochem Biophys* 2009;483:127–  
307 135.
- 308 [14] Jung T, Engels M, Kaiser B, Poppek D, Grune T. Intracellular distribution of oxidized  
309 proteins and proteasome in HT22 cells during oxidative stress. *Free Radic Biol Med*  
310 2006;40:1303–1312.
- 311 [15] Bloomer RJ, Goldfarb AH, Wideman L, McKenzie MJ, Consitt LA. Effects of acute  
312 aerobic and anaerobic exercise on blood markers of oxidative stress. *J Strength Cond*  
313 *Res* 2005;19:276–85.
- 314 [16] Chevion S, Moran DS, Heled Y, Shani Y, Regev G, Abbou B, Berenshtein E,  
315 Stadtman ER, Epstein Y. Plasma antioxidant status and cell injury after severe  
316 physical exercise. *Proc Natl Acad Sci U S A* 2003;100:5119–23.
- 317 [17] McGinnis G, Kliszczewicz B, Barberio M, Ballmann C, Peters B, Slivka D, Dumke  
318 C, Cuddy J, Hailes W, Quindry J. Acute Hypoxia and Exercise-Induced Blood  
319 Oxidative Stress. *Int J Sport Nutr Exerc Metab* 2014;
- 320 [18] Guidi F, Magherini F, Gamberi T, Bini L, Puglia M, Riccardo Marzocchi FR,  
321 Amedeo Modesti MGAM Pietro. Plasma protein carbonylation and physical exercise.  
322 *Mol BioSyst* 2011;7:640–650.
- 323 [19] Dalle-Donne I, Rossi R, Giustarini D, Milzani A, Colombo R. Protein carbonyl groups  
324 as biomarkers of oxidative stress. *Clin Chim Acta* 2003;329:23–38.

- 325 [20] Jung T, Höhn A, Grune T. The proteasome and the degradation of oxidized proteins:  
326 Part II – protein oxidation and proteasomal degradation. *Redox Biol* 2014;2:99–104.
- 327 [21] Jung T, Grune T. The proteasome and its role in the degradation of oxidized proteins.  
328 *IUBMB Life* 2008;60:743–752.
- 329 [22] Dalle-Donne I, Aldini G, Carini M, Colombo R, Rossi R, Milzani A. Protein  
330 carbonylation, cellular dysfunction, and disease progression. *J Cell Mol Med*  
331 2006;10:389–406.
- 332 [23] Jackson MJ. Redox regulation of adaptive responses in skeletal muscle to contractile  
333 activity. *Free Radic Biol Med* 2009;47:1267–1275.
- 334 [24] Babior BM. Phagocytes and oxidative stress. *Am J Med* 2000;109:33–44.
- 335 [25] Li JM, Shah AM. Endothelial cell superoxide generation: regulation and relevance for  
336 cardiovascular pathophysiology. *Am J Physiol Integr Comp Physiol* 2004;287:R1014–  
337 R1030.
- 338 [26] Balon TW, Nadler JL. Nitric oxide release is present from incubated skeletal muscle  
339 preparations. *J Appl Physiol* 1994;77:2519–2521.
- 340 [27] Mcardle A, Pattwell D, Vasilaki A, Griffiths RD, Jackson MJ, Physiol AJ, Metab E.  
341 Contractile activity-induced oxidative stress : cellular origin and adaptive responses  
342 Contractile activity-induced oxidative stress : cellular origin and adaptive responses.  
343 *Society* 2010;621–627.
- 344 [28] O'Neill CA, Stebbins CL, Bonigut S, Halliwell B, Longhurst JC. Production of  
345 hydroxyl radicals in contracting skeletal muscle of cats. *J Appl Physiol* 1996;81:1197–  
346 1206.
- 347 [29] Winterbourn CC. The biological chemistry of hydrogen peroxide. 1st ed. Elsevier Inc.;  
348 2013.
- 349 [30] Lamprecht M, Greilberger JF, Schwabegger G, Hofmann P, Oettl K. Single bouts of  
350 exercise affect albumin redox state and carbonyl groups on plasma protein of trained  
351 men in a workload-dependent manner. *J Appl Physiol* 2008;104:1611–1617.
- 352 [31] Michailidis Y, Jamurtas AZ, Nikolaidis MG, Fatouros IG, Koutedakis Y, Papassotiriou  
353 I, Kouretas D. Sampling time is crucial for measurement of aerobic exercise-induced  
354 oxidative stress. *Med Sci Sports Exerc* 2007;39:1107–13.
- 355 [32] Grune T, Reinheckel T, Davies KJA. Degradation of oxidized proteins in mammalian.  
356 *Faseb J* 1997;11:526–534.
- 357 [33] Pantke U, Volk T, Schmutzler M, Kox WJ, Sitte N, Grune T. Oxidized proteins as a  
358 marker of oxidative stress during coronary heart surgery. *Free Radic Biol Med*  
359 1999;27:1080–1086.
- 360 [34] Sachdev S, Davies KJA. Production, detection, and adaptive responses to free radicals  
361 in exercise. *Free Radic Biol Med* 2008;44:215–223.
- 362 [35] Gochman E, Reznick A., Avizohar O, Ben-Amotz A, Levy Y. Exhaustive exercise  
363 modifies oxidative stress in smoking subjects. *Am J Med Sci* 2007;333:346–353.
- 364 [36] Nikolaidis M., Jamurtas A., Paschalis V, Kostaropoulos I., Kladi-Skandali A,  
365 Balamitsi V, Koutedakis Y, Kouretas D. Exercise-induced oxidative stress in G6PD-  
366 deficient individuals. *Med Sci Sport Exerc* 2006;38:1443–1450.



- 367 [37] Bloomer RJ, Fry AC, Falvo MJ, Moore CA. Protein carbonyls are acutely elevated  
368 following single set anaerobic exercise in resistance trained men. *J Sci Med Sport*  
369 2007;10:411–417.
- 370 [38] Turner JE, Hodges NJ, Bosch JA, Aldred S. Prolonged Depletion of Antioxidant  
371 Capacity after Ultraendurance Exercise. *Med Sci Sport Exerc* 2011;43:1770–1776.
- 372 [39] Khassaf M, Child RB, McArdle A, Brodie DA, Esanu C, Jackson MJ. Time course of  
373 responses of human skeletal muscle to oxidative stress induced by nondamaging  
374 exercise. *J Appl Physiol* 2001;90:1031–1035.
- 375 [40] Khassaf M, McArdle A, Esanu C, Vasillaki A, McArdle F, Griffiths R., Brodie D.,  
376 Jackson M. Effect of vitamin C supplements on antioxidant defence and stress proteins  
377 in human lymphocytes and skeletal muscle. *J Physiol* 2003;549:645–652.
- 378 [41] Ji LL. Modulation of skeletal muscle antioxidant defense by exercise: Role of redox  
379 signaling. *Free Radic Biol Med* 2008;44:142–52.
- 380 [42] Abed KE, Rebai H, Bloomer RJ, Trabelsi K, Masmoudi L, Zbidi A, Sahnoun Z,  
381 Hakim A, Tabka Z. Antioxidant status and oxidative stress at rest and in response to  
382 acute exercise in judokas and sedentary. *J Strength Cond Res* 2004;25:2400–2409.
- 383 [43] Miyazaki H, Oh-ishi S, Ookawara T, Kizaki T, Toshinai K, Ha S, Haga S, Ji LL, Ohno  
384 H. Strenuous endurance training in humans reduces oxidative stress following  
385 exhausting exercise. *Eur J Appl Physiol* 2001;84:1–6.
- 386 [44] Radovanovic D, Bratic M, Nurkic M, Cvetkovic T, Ignjatovic A, Aleksandrovic M.  
387 Oxidative stress biomarker response to concurrent strength and endurance training.  
388 *Gen Physiol Biophys* 2009;28:205–211.
- 389 [45] Fisher G, Pascoe DD, Schwartz DD, Quindry J, Barberio MD, Foster EB, Jones KW.  
390 Lymphocyte enzymatic antioxidant responses to oxidative stress following high-  
391 intensity interval exercise Lymphocyte enzymatic antioxidant responses to oxidative  
392 stress following high-intensity interval exercise. *J Appl Physiol* 2011;110:730–737.
- 393 [46] Shi M, Wang X, Yamanaka T, Ogita F, Nakatani K, Takeuchi T. Effects of anaerobic  
394 exercise and aerobic exercise on biomarkers of oxidative stress. *Environ Health Prev*  
395 *Med* 2007;12:202–208.
- 396 [47] Rahnema N, Gaeini AA, Hamedinia MR. Oxidative stress responses in physical  
397 education students during 8 weeks aerobic training. *J Sport Med Phys Fitness*  
398 2007;47:119–123.
- 399 [48] Kabasakalis, A., Kyparos, A., Tsalis, G., Loupos, D., Pavlidou, A., and Kouretas D.  
400 Blood oxidative stress markers after ultramarathon swimming. *J Strength Cond Res*  
401 2011;25:805–811.
- 402 [49] Gaeini A., Rahnema N, Hamedinia M. Effects of vitamin E supplementation on  
403 oxidative stress at rest and after exercise to exhaustion in athletic students. *J Sport Med*  
404 *Phys Fitness* 2006;46:458–461.
- 405 [50] Magalhães J, Ferreira R, Marques F, Olivera E, Soares J, Ascensão A. Indoor climbing  
406 elicits plasma oxidative stress. *Med Sci Sport Exerc* 2007;39:955–963.
- 407 [51] Morillas-Ruiz J, Zafrilla P, Almar M, Cuevas MJ, López FJ, Abellán P, Villegas J a,  
408 González-Gallego J. The effects of an antioxidant-supplemented beverage on exercise-  
409 induced oxidative stress: results from a placebo-controlled double-blind study in  
410 cyclists. *Eur J Appl Physiol* 2005;95:543–9.

- 411 [52] Wadley AJ, Chen YW, Bennett SJ, Lip GYH, Turner JE, Fisher J., Aldred S.  
412 Monitoring changes in thioredoxin and over-oxidised peroxiredoxin in response to  
413 exercise in humans. *Free Radic Res* 2015;49:290–298.
- 414 [53] Sitte N, Merker K, Grune T. Proteasome-dependent degradation of oxidized proteins  
415 in MRC-5 fibroblasts. *FEBS Lett* 1998;440:399–402.
- 416 [54] Egerer K, Kuckelkorn U, Rudolph PE, Rückert JC, Dörner T, Burmester G-R, Kloetzel  
417 P-M, Feist E. Circulating proteasomes are markers of cell damage and immunologic  
418 activity in autoimmune diseases. *J Rheumatol* 2002;29:2045–2052.
- 419 [55] Wada M, Kosaka M, Saito S, Sano T, Tanaka K, Ichihara A. Serum concentration and  
420 localization in tumor cells of proteasomes in patients with hematologic malignancy  
421 and their pathophysiologic significance. *Lab Clin Med* 1993;121:215–223.
- 422 [56] Hoffmann O, Heubner M, Anlasik T, Winterhalter M, Dahlmann B, Kasimir-Bauer S,  
423 Kimmig R, Wohlschlaeger J, Sixt SU. Circulating 20S proteasome in patients with  
424 non-metastasized breast cancer. *Anticancer Res* 2011;31:2197–2201.
- 425 [57] Zoeger A, Blau M, Egerer K, Feist E, Dahlmann B. Circulating proteasomes are  
426 functional and have a subtype pattern distinct from 20S proteasomes in major blood  
427 cells. *Clin Chem* 2006;52:2079–2086.
- 428 [58] Nedergaard A, Vissing K, Overgaard K, Kjaer M, Schjerling P. Expression patterns of  
429 atrogenic and ubiquitin proteasome component genes with exercise: effect of different  
430 loading patterns and repeated exercise bouts. *J Appl Physiol* 2007;103:1513–1522.
- 431 [59] Dalbo V, Roberts M., Hassell S, Kerksick C. Effects of pre-exercise feeding on serum  
432 hormone concentrations and biomarkers of myostatin and ubiquitin proteasome  
433 pathway activity. *Eur J Nutr* 2013;52:477–487.
- 434 [60] Urso M., Clarkson P., Hittel D, Hoffman E., Thompson P. Changes in ubiquitin  
435 proteasome pathway gene expression in skeletal muscle with exercise and statins.  
436 *Arter Thromb Vasc Biol* 2005;25:2560–2566.
- 437 [61] Jamart C, Francaux M, Millet GY, Deldicque L, Frère D, Féasson L. Modulation of  
438 autophagy and ubiquitin-proteasome pathways during ultra-endurance running. *J Appl*  
439 *Physiol* 2012;112:1529–1537.
- 440 [62] Ogonovszky H, Berkes I, Kumagai S, Kaneko T, Tahara S, Goto S, Radák Z. The  
441 effects of moderate-, strenuous- and over-training on oxidative stress markers, DNA  
442 repair, and memory, in rat brain. *Neurochem Int* 2005;46:635–640.
- 443 [63] Radak Z, Chung HY, Goto S. Exercise and hormesis: oxidative stress-related  
444 adaptation for successful aging. *Biogerontology* 2005;6:71–5.
- 445 [64] Augustyniak E, Adam A, Wojdyla K, Rogowska-Wrzesinska A, Willetts R, Korkmaz  
446 A, Atalay M, Weber D, Grune T, Borsa C, Gradinaru D, Chand Bollineni R, Fedorova  
447 M, Griffiths HR. Validation of protein carbonyl measurement: A multi-centre study.  
448 *Redox Biol* 2015;4:149–157.
- 449 [65] Madian AG, Regnier FE. Profiling Carbonylated Proteins in Human Plasma. *J*  
450 *Proteome Res* 2010;9:1330–1343.
- 451 [66] Montelpare WJ, Klentrou P, Thoden J. Continuous versus intermittent exercise effects  
452 on urinary excretion of albumin and total protein. *J Sci Med Sport* 2002;5:219–28.
- 453 [67] Junglee NA, Lemmey AB, Burton M, Searrell C, Jones D, Lawley JS, Jibani MM,  
454 Macdonald JH. Does Proteinuria-Inducing Physical Activity Increase Biomarkers of  
455 Acute Kidney Injury? *Kidney Blood Press Res* 2012;278–289.

- [68] Forslund A., Hambraeus L, Beurden H van, Holmbäck U, El-Khoury A., Hjorth G, Olsson R, Stridsberg M, Wide L, Akerfeldt T, Regan M, Young VR. Inverse relationship between protein intake and plasma free amino acids in healthy men at physical exercise. *Am J Physiol Endocrinol Metab* 2000;278:E857–E867.
- [69] Sheffield-Moore M, Paddon-Jones D, Sanford AP, Rosenblatt JJ, Matlock AG, Cree MG, Wolfe RR. Mixed muscle and hepatic derived plasma protein metabolism is differentially regulated in older and younger men following resistance exercise. *Am J Physiol Endocrinol Metab* 2005;288:E922–E929.
- [70] Michailidis Y, Jamurtas AZ, Nikolaidis MG, Fatouros IG, Koutedakis Y, Papassotiriou I, Kouretas D. Sampling time is crucial for measurement of aerobic exercise-induced oxidative stress. *Med Sci Sport Exerc* 2007;39:1107–1113.
- [71] Dean RT, Fu S, Stocker R, Davies MJ. Biochemistry and pathology of radical-mediated protein oxidation. *Biochem J* 1997;324 ( Pt 1:1–18.
- [72] Mohanty JG, Bhamidipaty S, Evans MK, Rifkind JM. A fluorimetric semi-microplate format assay of protein carbonyls in blood plasma. *Anal Biochem* 2010;400:289–94.
- [73] Alessio HM, Hagerman AE, Fulkerson BK, Ambrose J, Rice RE, Wiley RL. Generation of reactive oxygen species after exhaustive aerobic and isometric exercise. *Med Sci Sports Exerc* 2000;32:1576–1581.
- [74] Nikolaidis M., Kyparos A, Hadziioannou M, Panou N, Samaras L, Jamurtas A., Kouretas D. Acute exercise markedly increases blood oxidative stress in boys and girls. *Appl Physiol Nutr Metab* 2007;32:197–205.
- [75] Bloomer RJ, Creasy AK, Smith WA. Physical work-induced oxidative stress is exacerbated in young cigarette smokers. *Nicotine Tob Res* 2007;9:205–211.
- [76] Goldfarb A., McKenzie M., Bloomer R. Gender comparisons of exercise-induced oxidative stress: influence of antioxidant supplementation. *Appl Physiol Nutr Metab* 2007;32:1124–1231.
- [77] Bloomer R., Goldfarbm A., McKenzie M. Oxidative stress response to aerobic exercise: comparison of antioxidant supplements. *Med Sci Sport Exerc* 2006 Jun;38(6):1098-105 2006;38:1098–1105.
- [78] Goldfarb A., Patrick S., Bryer S, You T. Vitamin C supplementation affects oxidative-stress blood markers in response to a 30-minute run at 75% VO<sub>2</sub>max. *Int J Sport Nutr Exerc Metab* 2005;15:279–90.

## Figure Legends

**Figure 1: Formation of non-native carbonyl groups in proteins.** Non-native carbonyl groups can be introduced into proteins by; (1) direct oxidation of amino acids, (2) via secondary oxidation products of lipid peroxidation, and (3) the oxidation products of reducing sugars. An example carbonyl modification is included for each pathway; (A) (2-amino-3-ketobutyric acid, (B) malondialdehyde-Lysine adduct, and (C) carboxymethyl lysine (3)). The non-native carbonyl group in each example is indicated by a dashed circle around the C=O bond of the amino acid or amino acid side chain (adduct).

**Figure 2: Published and unpublished data indicating decreases in protein carbonyl concentration (nM/mg protein) in response to exercise.** PC level decreases during the last minute (End-Exercise) (A) 27 minutes of cycling at 60%  $\dot{V}O_{2\max}$  and (B) 20 minutes of cycling at 80%  $\dot{V}O_{2\max}$  in untrained young men (n=10; mean  $\pm$  SD: age  $22 \pm 3$  yrs;  $\dot{V}O_{2\max}$   $42.7 \pm 5.0$  ml/kg/min) [8]; C) PC level decreases immediately following a  $\dot{V}O_{2\max}$  test to exhaustion in trained young men (n=10; mean  $\pm$  SD: age  $23 \pm 3$  yrs;  $\dot{V}O_{2\max}$   $61.9 \pm 4.7$  ml/kg/min) (Turner & Aldred; unpublished work); D) PC level decreases immediately following 75 minutes of cycling at 70%  $\dot{V}O_{2\max}$ , and is elevated above baseline levels 2 hours post-exercise in trained young men (n=12; mean  $\pm$  SD: age:  $28 \pm 4$  years,  $\dot{V}O_{2\max}$   $63.7 \pm 5.3$  ml/kg/min) (Wadley et al, 2015; *in preparation*). Individual (grey bars) and mean (black line) PC concentration changes are reported. Mean percentage change and statistical significance are indicated above each graph. Paired samples T-tests (A-C) and repeated measures ANOVA (D) were performed using SPSS (PASW Statistics, 22.0).

**Figure 3: Proposed formation/ clearance of protein carbonyl groups in response to exercise in humans.** Exercise can cause an increase in plasma protein carbonylation when the exercise bout is of sufficient intensity and/or duration. Activation of clearance mechanisms may drive a decrease in PC groups present at baseline when certain exercise conditions are met.